Summary of Group Mean Maternal and Fetal Observations at Cesarean Section

	Sodium Hippurate (mg/kg/day)					
	1118			5590		
	No.	%	S.D.	No.	%	S.D.
Animals on study:	25	-	-	25	-	-
Animals that were gravid:	24	96.0	-	25	100.0	-
Animals that died:	0	0.0	-	11	44.0	-
Nongravid:	0	0.0	-	0	0.0	-
Gravid:	0	0.0	-	11	100.0	-
Animals that aborted/delivered:	0	0.0	-	0	0.0	-
Animals examined at Cesarean Section:	25	100.0	-	14	56.0	-
Nongravid:	1	4.0	-	0	0.0	-
Gravid:	24	96.0	-	14	100.0	-
Dams with resorptions only:	0	0.0	-	1	7.1	-
Dams with viable fetuses:	24	100.0	-	13	92.9	-
Viable fetuses/dam:	13.9	-	±2.58	12.6	-	±5.29
Postimplantation loss/dam:	1.3	-	±2.05	2.2	-	±3.70
Total Implantations/dam:	15.1	-	±1.39	14.9	-	±3.96
Corpora lutea/dam:	16.6	-	±2.00	17.9	-	±3.55
Group mean preimplantation loss (%) [a]:	-	9.0	-	-	16.8	-
Group mean preimplantation loss (%) [b]:	-	8.3	-	-	14.9	-
Fetal sex distribution – male:	173	52.0	-	94	53.1	-
- female:	160	48.0	-	83	48.9	-
Mean fetal body weight (grams):	3.7	-	±0.40	3.0[c,d]	-	±0.61
Mean crown rump length (cm):	3.8	-	±0.16	3.5	-	±0.24

[a] $\frac{\text{Total No. Corpora Lutea} - \text{Total No. Implantations}}{\text{Total No. Corpora Lutea}} \times 100 = \text{Group Mean Preimplantation Loss (\%)}$

[b] $\frac{\text{Total No. Implantations} - \text{Total No. Viable Fetuses}}{\text{Total No. Implantations}} \times 100 = \text{Group Mean Preimplantation Loss (\%)}$

[c] Significantly different from the vehicle control group mean; $p < 0.01$

[d] Significantly different from the untreated control group mean; $p < 0.01$

[e] Value does not include dam #92289 with an erroneous number of corpora lutea

S.D. – Standard deviation

- Not applicable