Table 2: Percent incidence of skeletal anomalies among offspring of mice treated with dinoseb (gestational days 10, 11, & 12) after food deprivation (gestational days 9 or 9 & 10)

	Dose (mg/kg/day)								
	0			14.1			15.8		
Type of anomaly	0 (14)	24 (14)	48 (7)	0 (7)	24 (6)	48 (6)	0 (15)	24 (14)	48 (6)
Digital bones (FL)[b] abs or NO[c]	0[d]	1.2 (1.2)	0	0	5.6 (5.6)	20.0 (16.3)*†[e]	8.5 (3.1)	18.4 (8.0)*	6.9 (3.1)
Ischium or pubis abs or small	0	1.0 (1.0)	0	2.0 (2.0)	11.1 (11.1)	8.3 (8.3)	27.6 (9.0)*	27.6 (7.1)*	14.0 (5.2)
Femur abs or small	1.4 (1.4)	0	0	2.0 (2.0)	18.0 (13.7)	12.5 (12.5)	19.5 (7.3)*	40.2 (9.8)*†	14.0 (5.2)
Tibia or fibula abs or small	0	0	0	0	18.0 (13.7)	12.5 (12.5)	20.7 (8.2)*	46.2 (10.1)*†	16.4 (5.8)
Digital bones (HL)[f] abs or NO	0	1.2 (1.2)	0	0	12.5 (8.5)	15.8 (12.30	11.4 (5.4)	40.1 (9.4)*†	9.5 (6.0)
Phalanges (HL) abs or NO	18.5 (7.1)	23.3 (7.5)	40.2 (13.0)	37.2 (12.9)	43.2 (18.3)	74.4 (14.0)†	32.3 (6.9)	58.9 (9.9)*†	54.8 (14.4)
Hallux abs or NO	27.0 (8.7)	38.4 (9.3)	70.8 (11.4)†	84.9 (8.9)*	77.9 (11.4)*	97.6 (2.4)	79.8 (5.6)*	80.2 (8.0)*	91.7 (8.3)
Ribs fused	0	8.3 (5.7)	12.2 (12.2)	8.2 (5.3)	5.2 (3.3)	14.0 (8.0)	17.6 (5.6)*	35.7 (7.4)*†	14.3 (9.0)
Vertebrae incomplete	2.4 (2.4)	2.1 (1.4)	16.3 (14.1)	10.8 (4.3)	11.1 (11.1)	29.8 (15.0)	26.3 (6.8)*	37.0 (8.9)*	12.3 (6.8)
Vertebrae fused	0	3.6 (3.6)	6.1 (6.1)	23.0 (12.0)	33.1 (15.0)*	39.4 (14.6)*	59.6 (9.3)*	63.9 (9.9)*	61.4 (10.5)*
Vertebrae split	0	5.7 (3.7)	0	0	2.4 (2.4)	0	17.0 (4.0)*	14.3 (4.7)	11.0 (7.1)
Sternabrae abs or NO	0	2.0 (1.4)	0	0	25.4 (14.0)*†	10.7 (8.2)	18.0 (4.1)*	42.0 (4.8)*†	13.1 (7.9)
Sternabrae split	6.0 (2.9)	20.5 (6.9)	18.2 (8.0)	21.3 (9.2)	2.4 (2.4)	30.2 (8.2)	30.6 (5.6)*	23.7 (4.8)	39.6 (7.8)

[a] The number in parentheses given with the deprivation time is the number of litters examined.
[b] FL = front limb
[c] abs or NO = absent or not ossified
[d] Values are percent incidence mean response/litter (SE)
[e] Values marked with an asterisk (*) differ significantly from those of the similarly deprived, no drug group; $p<0.05$; values marked with a dagger (†) differ significantly from those of the non-deprived group that was treated with the same dose of test material; $p<0.05$
[f] HL = hind limb