

Table 2

Body Weights of Mice receiving Ionox 330 for two years

	Dietary Conc (ppm)	Body Weight (g) at Week								
		0	13	26	39	52	64	78	92	104
Males										
	0	23.8	31.7	34.9	39.3	39.4	39.9	40.8	42.0	36.1
	1000	24.5	31.9	35.3	39.9	39.8	41.4	42.3	43.7	35.8
	5000	24.0	32.0	36.3	39.6	39.8	40.8	41.8	43.5	
	Standard error of treatment Mean	± 0.52	± 0.45	±0.47	± 0.51	± 0.51	± 0.61	± 0.77	± 0.68	± 1.13
		± 0.57	± 0.49	± 0.51	± 0.55	± 0.55	± 0.67	± 0.83	± 0.80	± 1.35
							± 0.68	± 0.88	± 0.81	± 1.32
Females										
	0	20.2	28.6	32.3	33.6	33.9	35.0	34.6	35.5	32.9
	1000	19.7	27.6	31.8	33.7	34.5	34.8	35.7	35.9	33.7
	5000	19.6	27.8	30.8	32.7	33.2	33.3	33.2	35.4	30.3
	Standard error of treatment Mean	± 0.29	± 0.36	± 0.45	± 0.54	± 0.52	± 0.73	± 0.78	± 0.93	± 1.03
		± 0.31	±0.40	±0.51	± 0.63	± 0.59	± 0.83	± 0.90	± 1.02	± 1.17
				± 0.50	±0.60	± 0.57	± 0.80	± 0.84		± 1.12

Treatment means adjusted for differences in the initial body weight.
 ≤ 0.05 Significance of the difference between treatment and control mean